

Breast Cancer Screening (BCS-E)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Female patients ages 50-74 who had a mammogram screening completed</p> <p>TIMELINE On or by Oct. 1, two years prior to the measurement year through Dec 31 of the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Order/completion of Mammogram ✓ Document completion and result of mammogram in visit note 	<ul style="list-style-type: none"> ✓ Consultation reports ✓ Diagnostic reports ✓ Health history documentation 	<p>CPT: 77061, 77062, 77063, 77065, 77066, 77067</p> <p>CPT II: N/A</p>

Colorectal Cancer Screening (COL-E)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION 45-75 who had a screening for colorectal cancer</p> <p>TIMELINE - Colonoscopy (<i>measurement year or 9 years prior</i>) - Flexible sigmoidoscopy or CT colonography (<i>measurement year or 4 years prior</i>) - FIT-DNA test (<i>measurement year or 2 years prior</i>) - FOBT, gFOBT, FIT (<i>measurement year</i>)</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Order/completion of appropriate screening test ✓ Provide iFOBT or FIT-DNA test during visit ✓ Document completion and result of screening test in visit note 	<ul style="list-style-type: none"> ✓ Consultation reports ✓ Diagnostic reports ✓ Health history documentation ✓ Lab reports 	<p>CPT: 44388-44394, 44401-44408, 45378-45393, 45398</p> <p>CPT II: N/A</p>

Quality Reference Guide 2025

Diabetic Screenings



Eye Exam for Patients with Diabetes (EED)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18-75 with diabetes (Type 1 or 2) who had a diabetic retinal eye exam</p> <p>TIMELINE - Positive retinal or dilated eye exam (<i>measurement year</i>) - Negative retinal or dilated eye exam (<i>measurement year or one year prior</i>) - Bilateral eye enucleations (<i>anytime in patient's history through Dec. 31 of measurement year</i>)</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Refer to Optometrist or Ophthalmologist for test ✓ Document completion and result of screening test in visit note 	<ul style="list-style-type: none"> ✓ Consultation reports ✓ Diabetic flow sheets ✓ Progress notes ✓ Eye exam report note 	<p>CPT: 67028, 67030, 67031, 67036, 67039, 67040, 67041-3, 67101, 67105, 67107, 67108, 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220, 67221, 67227, 67228, 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92227, 92228, 92230, 92235, 92240, 92250, 92260</p> <p>CPT II: 2023F, 2025F, 2033F, 2022F, 2024F, 2026F, 3072F (Billed by Optometrist or Ophthalmologist)</p>

Glycemic Status Assessment for Patients With Diabetes (GSD)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18-75 with diabetes (Type 1 or 2) who had an HBA1c test during the measurement year that showed an A1C under control ($\leq 9.0\%$) *Glucose Management Indicator results included</p> <p>TIMELINE Anytime during the measurement year (<i>if multiple performed, the last test result is used</i>)</p> <p>CMS WEIGHT: 3</p>	<ul style="list-style-type: none"> ✓ Order/complete test ✓ Write extended-day prescriptions (90-100 day) 	<ul style="list-style-type: none"> ✓ Consultation reports ✓ Diabetic flow sheets ✓ Progress notes ✓ Lab reports ✓ Vital sheets (all documentation needs date of test and result) 	<p>CPT: N/A</p> <p>CPT II: 3044F, 3051F, 3052F, 3046F</p>

Kidney Health Evaluation for Patients with Diabetes (KED)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18-85 with diabetes (Types 1 or 2) who had a kidney health evaluation (both an eGFR and a uACR test are required)</p> <p>TIMELINE Anytime during the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Order and complete eGFR AND ✓ 1 urine albumin-creatinine ratio test (uACR) or urine albumin and urine creatinine tests for uACR within 4 days of each other 	<p>N/A: This measure can only be closed via claims</p>	<p>CPT: 80047, 80048, 80050, 80053, 80069, 82565, 82043, 82570</p> <p>CPT II: N/A</p>

Medication Adherence for Cholesterol (MAC)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18 or older who are compliant with their cholesterol medication (statin) at least 80% of the time in the measurement period</p> <p>TIMELINE Patients qualify for measure after the second fill of the drug, but the measurement period starts with the date of the first fill</p> <p>CMS WEIGHT: 3</p>	<ul style="list-style-type: none"> ✓ Consider writing a 3-month supply of prescription and offer a 100-day refill if available ✓ Discuss home delivery prescription options ✓ Ensure prescription reflects the current dosage ✓ Discuss any barriers: side effects, cost and transportation 	<p>N/A: This measure can only be closed via pharmacy claims</p>	<p>N/A: This measure can only be closed via pharmacy claims for refills</p>

Medication Adherence for Diabetes (MAD)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18 or older who are compliant with their diabetic medication at least 80% of the time in the measurement period (<i>excludes insulin</i>)</p> <p>TIMELINE Patients qualify for measure after the second fill of the drug, but the measurement period starts with the date of the first fill</p> <p>CMS WEIGHT: 3</p>	<ul style="list-style-type: none"> ✓ Consider writing a 3-month supply of prescription and offer a 100-day refill if available ✓ Discuss home delivery prescription options ✓ Ensure prescription reflects the current dosage ✓ Discuss any barriers: side effects, cost and transportation 	<p>N/A: This measure can only be closed via pharmacy claims</p>	<p>N/A: This measure can only be closed via pharmacy claims for refills</p>

Medication Adherence for Hypertension (MAH)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18 or older who are compliant with their hypertension medication (RAS antagonist) at least 80% of the time in the measurement period</p> <p>TIMELINE Patients qualify for measure after the second fill of the drug, but the measurement period starts with the date of the first fill</p> <p>CMS WEIGHT: 3</p>	<ul style="list-style-type: none"> ✓ Consider writing a 3-month supply of prescription and offer a 100-day refill if available ✓ Discuss home delivery prescription options ✓ Ensure prescription reflects the current dosage ✓ Discuss any barriers: side effects, cost and transportation 	<p>N/A: This measure can only be closed via pharmacy claims</p>	<p>N/A: This measure can only be closed via pharmacy claims for refills</p>

Statin Therapy for Patients with Cardiovascular Disease (SPC)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Male patients ages 21-75 and female patients 40-75 who have atherosclerotic cardiovascular disease (ASCVD) and receive at least 1 high or moderate intensity statin medication</p> <p>TIMELINE Measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Consider prescribing a statin as appropriate ✓ Recommend automatic refills to support adherence 	<p>N/A: This measure can only be closed via claims</p>	<p>N/A: This measure can only be closed via pharmacy claims for refills</p>

Statin Use in Persons with Diabetes (SUPD)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Diabetic patients ages 40-75 who have received at least 1 fill of a statin medication (<i>Patients with diabetes are defined as those who have had 2 fills of a diabetic medication during the measurement year</i>)</p> <p>TIMELINE Measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Consider prescribing a statin as appropriate ✓ Recommend automatic refills to support adherence 	<p>N/A: This measure can only be closed via claims</p>	<p>N/A: This measure can only be closed via pharmacy claims for refills</p>

Controlling Blood Pressure (CBP)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18-85 who had a diagnosis of hypertension and whose blood pressure was controlled at <140/90mmHg</p> <p>TIMELINE Anytime during the measurement year (if multiple performed, the last test result is used)</p> <p>CMS WEIGHT: 3</p>	<ul style="list-style-type: none"> ✓ Advise patients not to smoke, drink caffeinated drinks or exercise 30 minutes before BP reading ✓ Document each BP reading at each visit and if initial reading is >140/90 repeat before the patient leaves the office 	<ul style="list-style-type: none"> ✓ Consult notes ✓ Progress notes ✓ Medical history ✓ Vital sheets 	<p>CPT: N/A</p> <p>CPT II: 3074F, 3075F, 3077F, 3078F, 3079F, 3080F</p>

Osteoporosis Management in Women Who Had a Fracture (OMW)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Women ages 67-85 who had a fracture and had a bone mineral density (BMD) test or a prescription to treat osteoporosis within 6 months from the date of the fracture (does not include finger, toe, face or skull fractures)</p> <p>TIMELINE Within 6 months of the date of the fracture</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ See patients in the office soon after the fracture occurs ✓ Schedule and order BMD at the time of the office visit 	<ul style="list-style-type: none"> ✓ Progress notes ✓ Medication list 	<p>CPT: 76977, 77078, 77080, 77081, 77085, 77086</p> <p>CPT II: N/A</p>

Plan All-Cause Readmissions (PCR)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18 years and older with an acute inpatient and observation stays that were followed by an unplanned acute readmission for any diagnosis within 30 days</p> <p>TIMELINE Anytime during the measurement year</p> <p>CMS WEIGHT: 3</p>	<ul style="list-style-type: none"> ✓ Follow-up with patients within 1 week of discharge ✓ Create a post-discharge outreach program to ensure patients understand discharge instructions, medication changes, follow-up appts, etc. 	N/A	<p>CPT: N/A</p> <p>CPT II: N/A</p>

Follow-up After Emergency Dept Visit For People with High-Risk Multiple Chronic Conditions (FMC)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients 18 years or older with an emergency department (ED) visit who have multiple high-risk chronic conditions who had a follow-up service within 7 days of ED visit</p> <p>TIMELINE Within 7 days of ED visit</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Provide patients with alternative options to the ED including urgent care, telehealth or in-person office visit ✓ Remind patients to schedule follow-up office or telehealth visit within 7 days of ED visit 	N/A	<p>CPT: N/A</p> <p>CPT II: N/A</p>

Transition of Care - Patient Engagement After Inpatient Discharge (TRCPE)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 18 years and older with non-acute inpatient discharge with patient engagement documented</p> <p>TIMELINE Within 30 days of discharge</p> <p>CMS WEIGHT: 0</p>	<ul style="list-style-type: none"> ✓ Use a variety of methods to engage patient: outpatient office, telephone, virtual or transitional care management visits 	<ul style="list-style-type: none"> ✓ Progress notes ✓ Health history and physical ✓ Home health records 	<p>CPT: Outpatient visits, transitional care management codes</p> <p>CPT II: N/A</p>

Transition of Care - Medication Reconciliation Post-Discharge (TRCMRP)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients 18 years or older with an acute or non-acute inpatient discharge with medication reconciliation documented</p> <p>TIMELINE Day of discharge or 30 days after</p> <p>CMS WEIGHT: 0</p>	<ul style="list-style-type: none"> ✓ Schedule office visits as soon as possible with patient after discharge ✓ If patients are unable to attend an office visit, medication reviews may be completed virtually, telehealth, via email through a secure portal or email 	<ul style="list-style-type: none"> ✓ Progress notes ✓ Health history and physical ✓ Home health records ✓ Medication list 	<p>CPT: 99483, 99495, 99496</p> <p>CPT II: 1111F</p>

Care of Older Adults - Functional Status Assessment (COA)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 66 and older who had evidence of a functional status assessment</p> <p>TIMELINE In the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Clearly documented the assessment completed with results and date of service completed ✓ Assessment can be conducted via in person, phone, virtual, telehealth and not limited to Providers completing the assessment 	<ul style="list-style-type: none"> ✓ Progress notes ✓ Health history and physical ✓ Home health records ✓ Functional status forms 	<p>CPT: N/A</p> <p>CPT II: 1107F</p>

Care of Older Adults - Pain Assessment (COA)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 66 and older who were assessed for pain</p> <p>TIMELINE In the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Complete an annual pain screening with a standardized tool and document the results in the EMR 	<ul style="list-style-type: none"> ✓ Progress notes ✓ Health history and physical ✓ Home health records ✓ Pain assessment forms 	<p>CPT: N/A</p> <p>CPT II: 1125F, 1126F</p>

Care of Older Adults - Medication Review (COA)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Patients ages 66 and older who had a medication review by the prescribing provider or a clinical pharmacist AND the presence of a medication list in the medical record or transitional care management services</p> <p>TIMELINE In the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Clearly document the date of service of the completed review in the medical record or notation of no medications ✓ Review can be conducted via phone by the Prescribing Provider or Clinical Pharmacist 	<ul style="list-style-type: none"> ✓ Progress notes ✓ Health history and physical ✓ Medication list 	<p>CPT: N/A</p> <p>CPT II: 1159F with 1160F</p>

Concurrent Use of Opioids and Benzodiazepines (COB)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Percentage of patients 18 and older with concurrent use of prescription opioids and benzodiazepines</p> <p>TIMELINE Patients are in the numerator when at least 2 fills of an opioid and 2 fills of a benzodiazepine have an overlapping days' supply for at least 30 cumulative days during the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Review concurrent use and consider discontinuing one of the medications or use alternative therapies ✓ Assist patients in exploring alternative options for managing pain ✓ Coordinate care with the patient's entire care team to avoid co-prescriptions 	<p>N/A: Based on Pharmacy claims</p>	<p>N/A: Based on Pharmacy claims</p>

Polypharmacy-Use of Multiple Anticholinergics Medications in Older Adults (Poly-ACH)

Measure Requirements	Best Practices	Documentation Needed for Closure	Codes to Close Measure
<p>DESCRIPTION Percentage of patients 65 and older with concurrent use of 2 or more unique anticholinergic (ACH) medications</p> <p>TIMELINE Patients with at least 2 fills of each medication on different dates of service in the targeted drug class during the measurement year. Concurrent use is defined as overlapping days' supply for at least 30 cumulative days during the measurement year</p> <p>CMS WEIGHT: 1</p>	<ul style="list-style-type: none"> ✓ Educate patients on the risk of taking 2 or more anticholinergic medications ✓ Review therapies, duration and evaluate potential risk of continued therapy versus the benefits ✓ Discontinue medications as appropriate or consider safer alternatives 	<p>N/A: Based on Pharmacy claims</p>	<p>N/A: Based on Pharmacy claims</p>